

University of Groningen

[Poster on] The effects of classroom interventions on the behaviour of school-aged children with ADHD symptoms

Gaastra, Geraldina; Groen, Y.; Tucha, L.; Tucha, O.

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Early version, also known as pre-print

Publication date:

2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gaastra, G., Groen, Y., Tucha, L., & Tucha, O. (2014). *[Poster on] The effects of classroom interventions on the behaviour of school-aged children with ADHD symptoms: A meta-analysis*. Poster session presented at Eunethydis 3rd International Conference on ADHD, Istanbul, Turkey.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.


Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

THE EFFECTS OF CLASSROOM INTERVENTIONS ON THE BEHAVIOR OF SCHOOL-AGED CHILDREN WITH ADHD SYMPTOMS: A META-ANALYSIS

Gaastra, G.F., Groen, Y., Tucha, L., & Tucha, O.
Department of Clinical and Developmental Neuropsychology
University of Groningen, Groningen, The Netherlands



OBJECTIVE

Children with attention-deficit/hyperactivity disorder (ADHD) often exhibit problem behavior in class, whereas teachers often seem to lack the knowledge and skills to manage this behavior. The aim of the study was to determine the effectiveness of several types of classroom interventions that teachers can apply in order to decrease off-task and disruptive behavior of school-aged children with ADHD symptoms. Furthermore, potential moderators were examined.

METHOD

- Literature search in PsycINFO, ERIC, and Web of Science + reference lists of review studies and included studies
- Inclusion criteria
 - Study: Published in academic journal, English
 - Participants: ADHD symptoms, age 6 - 17 years
 - Intervention: Implementable by teacher
 - Outcome: On-task, off-task, disruptive behavior
- Separate meta-analyses for within-subjects (WSD) and single-subject design studies (SSD) using standardized mean differences
- Subgroup analyses for intervention type (antecedent-based, consequence-based, self-regulation, and combined) and other potential moderators (setting, age, gender, medication)

TYPES OF INTERVENTIONS

Antecedent-based

Manipulation environment, instruction, or task

Consequence-based

Reinforcement and punishment procedures

Self-regulation

Teaching self-control and problem-solving skills

RESULTS

	Effect size (standardized mean difference)	
	WSD (k = 23)	SSD ¹ (k = 68)
Overall	0.82**	2.90
Intervention type	Consequence-based > other**	
Antecedent-based	0.32** (k = 10)	2.63 (k = 15)
Consequence-based	1.71** (k = 7)	2.88 (k = 20)
Self-regulation	0.42 (k = 3)	3.18 (k = 24)
Combined	0.57* (k = 3)	2.56 (k = 9)
Setting	General education > other**	
General education	1.23** (k = 10)	3.19 (k = 33)
Other	0.48** (k = 13)	2.34 (k = 29)
Age	ns	
Children	0.89** (k = 21)	2.88 (k = 52)
Adolescents	0.26 (k = 2)	2.97 (k = 16)
Gender	ns	
0 – 19 % male	-	3.24 (k = 4)
20 – 80 % male	0.88** (k = 8)	3.14 (k = 8)
81 – 100 % male	0.82** (k = 14)	2.84 (k = 56)
Medication	ns	
0 – 19 % medicated	1.11** (k = 12)	2.86 (k = 18)
20 – 80 % medicated	0.19 (k = 2)	2.34 (k = 12)
81 – 100 % medicated	0.45 (k = 2)	3.09 (k = 22)

¹ No significance tests were performed for SSD studies
ns = not significant, * = p < 0.05, ** = p < 0.01

CONCLUSION

- Classroom interventions are effective in reducing off-task and disruptive behavior of children with ADHD symptoms
- Consequence-based interventions are most helpful, but SSD studies suggest also high effectiveness of self-regulation interventions
- Interventions in general education classrooms are more effective than those in other settings
→ More severe ADHD problems or already existing behavior programs in other settings?
- Age and medication do not influence effectiveness of classroom interventions
- No insight in impact of gender due to lack of studies that included females